DIRECTIONS FOR USE

Mixing Directions: Feed to heifers at the rate of 0.5 pounds per day. Mix with a complete ration. Continue feeding throughout the time heifers are being grown, finished and slaughtered.

For combination clearances, reference feed additive compendium or call Animal Science Product's technical service department.

INGREDIENTS

Guaranteed Analysis:

Melengestrol Acetate 0.00022% (Each pound contains 1.0mg Melengestrol Acetate)

Rice Hulls, Calcium carbonate, Mineral oil.

PACKAGING

50 lb. bag

Specially designed multi-wall bag contains 2 exterior walls of paper for protection, an inner-poly liner for moisture protection followed by and interior paper liner.

CAUTIONS

Discontinue feeding 48 hours before slaughter.

Store in a cool, dry place.

Keep out of reach of children.

Only for use in heifers being fed for slaughter. Not effective in spayed heifers and steers.

102894 MGA

MGA HEIFER SUPPLEMENT

TYPE "B" MEDICATED ARTICLE

ADVANTAGES

- For increased rate of gain.
- Improved feed efficiency.
- Suppression of estrus (heat) in heifers fed for slaughter.
- Contains AAFCO approved carrier.
- Production records and retained samples maintained for one year.
- Type "B" medicated article does not require 1900 registration.
- MSDS sheets available.

INDICATIONS

A type "B" medicated article specifically formulated for uniform dispersion in cattle feeds.